



NIH Public Access

Author Manuscript

J Cell Biochem. Author manuscript; available in PMC 2012 May 08.

Published in final edited form as:

J Cell Biochem. 2011 February ; 112(2): 607–613. doi:10.1002/jcb.22951.

Changes in Matrix Protein Gene Expression Associated With Mineralization in the Differentiating Chick Limb-bud Micromass Culture System

Cristina C. Teixeira¹, Jenny Xiang², Rani Roy³, Valery Kudrashov³, Itzhak Binderman^{3,4}, Philipp Mayer-Kuckuk³, and Adele L. Boskey^{2,3,*}

¹New York University, College of Dentistry, New York, New York

²Weill Medical College of Cornell University, New York, New York

³Mineralized Tissue Laboratory, Hospital for Special Surgery, New York, New York

⁴School of Dental Medicine, Tel Aviv University, Tel Aviv, Israel

Abstract

Chick limb-bud mesenchymal stem cells plated in high density culture in the presence of 4 mM inorganic phosphate and vitamin C differentiate and form a mineralizable matrix, resembling that of the chick growth plate. To further elucidate the mechanism that allows these cultures to form physiologic hydroxyapatite deposits, and how the process can be manipulated to gain insight into mineralization mechanisms, we compared gene expression in mineralizing (with 4 mM inorganic phosphate) and non-mineralizing cultures (containing only 1 mM inorganic phosphate) at the start of mineralization (day 11) and after mineralization reached a plateau (day 17) using a chick specific microarray. Based on replicate microarray experiments and K-cluster analysis, several genes associated with the mineralization process were identified, and their expression patterns confirmed throughout the culture period by quantitative RT-PCR. The functions of bone morphogenetic protein 1, BMP1, dentin matrix protein 1, DMP1, the sodium phosphate co-transporter, NaPi IIb, matrix metalloprotease 13, MMP-13, and alkaline phosphatase, along with matrix protein genes (type X collagen, bone sialoprotein, and osteopontin) usually associated with initiation of mineralization are discussed.

Keywords

CHICK LIMB-BUD; MICROMASS CULTURE; MINERALIZATION; MICROARRAY; GENE EXPRESSION

Chick limb-bud mesenchymal stem cells plated in high density (micromass) culture under previously defined conditions [Binderman et al., 1979; Boskey et al., 1991; Boskey et al., 1992b; Boskey et al., 1996], differentiate and form a mineralizable matrix, that resembles that of the chick epiphyseal growth plate [Boskey et al., 1992a]. This system has been used to study the effects of collagen [Boskey et al., 2000], proteoglycans [Boskey et al., 1997], and phosphoproteins [Boskey et al., 2008] on the mineralization process. To further elucidate the mechanism that allows these cultures to form physiologic hydroxyapatite deposits, and how the process can be manipulated to gain insight into mineralization

mechanisms, we compared gene expression in mineralizing and non-mineralizing cultures at the start of mineralization (day 11) and after mineralization reached a plateau (day 17) [Boskey et al., 2008] using a chick specific microarray.

While gene expression profiling is not generally reported for the epiphyseal growth plate mediated calcification, there are reports for gene expression in the early differentiation of mesenchymal cells into chondrocytes [Cameron et al., 2009], in the reserve [Zhang et al., 2008], proliferating and hypertrophic zones of the growth plate [Wang et al., 2004b], and in healing fracture callus development [Rundle et al., 2006; Wang et al., 2006; Khan et al., 2008] a process that mimics endochondral ossification. Similarly there are reports of gene expression during intramembranous bone formation [Kuroda et al., 2005], and for the effects of different factors on growth plate chondrocytes [Ulici et al., 2010]. There is also a proteomics study of the mouse growth plate [Belluccio et al., 2006] and of mineralizing extracellular matrix vesicles [Xiao et al., 2007]. Recently, James et al. [James et al., 2010] used microdissection of murine bones to characterize gene expression in a combination of proliferating and resting zone cells, prehypertrophic and hypertrophic chondrocytes, and calcified cartilage and bone and compared this data to micromass cultures of the murine cells. The emphasis in many of those studies was on signaling molecules and growth factors. The purpose of the present study was to identify genes associated with the mineralization process, and to select those extracellular matrix proteins and enzymes that modify them for further analysis.

MATERIALS AND METHODS

THE CULTURE SYSTEM

Chick limb-buds were obtained at stage 21–24 [Hamburger and Hamilton, 1951] from White Leg Horn chick embryos. Mesenchymal cells released by digestion in 0.05% trypsin-EDTA (Invitrogen) were separated from debris by passage through a 20 µm Nitex membrane, counted with a hemocytometer, checked for viability by trypan blue dye exclusion, and pelleted at 1100 rpm. Cells, resuspended in DME (Sloan Kettering Media Lab) containing 1 g/l glucose supplemented with 10% FBS (Invitrogen) and antibiotic/antimycotic (Invitrogen; 100 units penicillin, 100 µg streptomycin, 0.25 µg amphotericin B/ml), were plated using the micromass technique [Ahrens et al., 1977] at a density of 0.75×10^6 cells per 20 µl drop in 35mm Falcon tissue culture dishes. The micromass cultures were allowed to attach for 1.5 h in a humidified atmosphere of 5% CO₂ at 37°C, and then flooded with the same DME media. From day 2 onward, 25 µg/ml Vitamin C (Sigma) was added to the media along with 0.3 mg/ml L-glutamine (Invitrogen). At day 2, the calcium concentration of the media was adjusted to 1.3 mM using calcium chloride and the phosphate concentration was adjusted to 1.0 mM using monobasic sodium phosphate. For all mineralizing cultures, the total phosphate concentration (Pi) was adjusted to 4.0 mM with potassium acid phosphate (Fisher Chemicals, New Jersey). Control non-mineralizing cultures received no acid phosphate addition and were in 1.0 mM Pi. Cultures were incubated at 37° with 95% air, 5% CO₂ with media changed every 48 h. The overview of the experiment is presented schematically in Figure 1.

RNA ISOLATION

Following washing with phosphate buffered saline, total cellular RNA was isolated from mineralizing and non-mineralizing cultures on days 11 and 17 using the single step Qiagen RNeasy kit (Chatworth, CA), and used for gene expression profiling using the Affymetrix GeneChip chicken genome array containing 32,773 transcripts. Six cultures were used per time point and each culture system was repeated three times, providing triplicate RNA samples for analysis. The MessageAmp Premier RNA Amplification Kit (Ambion/Applied

Biosystems, Austin, TX), a one-round in vitro transcription (IVT) system was used to amplify nanogram amounts of total RNA from 6 samples and labeled cRNA for microarray hybridization. The quantity and quality of the amplified cRNA were assessed using NanoDrop ND-1000 Spectrophotometer (Thermo Scientific, Wilmington, DE) and Agilent Bioanalyzer (Santa Clara, CA). The biotin labeled cRNA was fragmented and hybridized to the Chicken Genome arrays (Santa Clara, CA). After Hybridization, GeneChip arrays were washed, stained, and scanned by GeneChip Scanner 3000 7G according to the Affymetrix Expression Analysis technical manual.

Affymetrix GeneChip Operating Software was used for image acquisition. The target signal intensity from each chip was scaled to 500. Analyses were done in the CUMC Genomics Resources Core Facility.

MICROARRAY ANALYSES

GeneChip chicken genome arrays (#900590) were acquired from Affymetrix (Affymetrix Inc., Santa Clara, CA), and were used according to the manufacturer's instructions. Three replicates of each experiment were analyzed for the data normalization, statistical analysis, and pattern study using GeneSpring software (Agilent, Redwood City, CA). Data was sorted for expression level based on p value ($P > 0.05$ was excluded) and fold increase/decrease. Data was plotted as K-clusters indicating genes that increased and or decreased in expression with time or with mineralization.

SELECTION OF GENES FOR qRT PCR

Genes were sorted into categories: signaling molecules and growth factors, genes associated with metabolism, inflammation, translation and transduction, matrix proteins and enzymes that modulate matrix proteins. Several of the latter were selected based on their postulated roles in the mineralization process, discussed later, for quantification by qRT PCR. Additionally we did qRT PCR on genes known to be involved in growth plate mineralization from earlier studies [Boskey et al., 1992b].

qRT PCR

RNA was isolated from the chick system at days 4, 7, 9, 11 or 12, 14, 16 and 21. Mineralizing (4P) and non-mineralizing cultures (1P) were included for each time point. In the first two experiments, collection was based on multiple batches of eggs. In the third and fourth experiment, the same batch of eggs and embryos provided all the cultures for all analysis. Expression of selected genes was analyzed by quantitative real-time qRT PCR using 5ng cDNA with the QuantiTect SYBR Green RT-PCR kit (Qiagen, Valencia, CA) on a DNA Engine Optican 2 System (MJ Research, Waltham, MA). Specific primers for these genes were obtained from Qiagen (Valencia, CA) using the chicken sequences with β -actin or GAPDH mRNA as a reference for quantification. The primers and their sources are summarized in Table I. Relative transcript levels are reported as "fold change" (x in the following formula) in gene expression and calculated using the threshold cycle (C_t) and the following formula, in which ctl =control, exp =experimental, and GAPDH or β -actin =the housekeeping gene: $x = 2^{\Delta\Delta C_t}$, where $\Delta\Delta C_t = \Delta E - \Delta C$, $\Delta E = C_t(\text{exp}) - C_t(\text{GAPDH})$, and $\Delta C = C_t(\text{ctl}) - C_t(\text{GAPDH})$. The change for $\Delta\Delta C_t$ was used in the calculation. Fold change was calculated relative to non-mineralizing controls.

STATISTICS

Mean and standard deviations for each group of genes analyzed in triplicate or quadruplicate were calculated (GraphPad, InStat) and compared to the 1.0 value indicating equivalence in

the culture. Prior to these comparisons, single outliers for fold change were excluded using the Q-test [Dean and Dixon, 1951]. Significance was assumed with $P < 0.05$.

RESULTS

Microarray analysis uncovered 1814 genes that were differentially expressed in mineralizing and control cultures at day 11 and 17. Only those genes that were elevated/suppressed more than two fold and had significant ($P < 0.05$) alterations in expression based on replicate microarrays from separate culture experiments were considered.

K-cluster analyses indicated genes that were significantly up or down regulated as mineralization progressed (Fig. 2) as opposed to those that were increased initially and then decreased or those that increased or decreased temporally rather than with mineralization. These genes had distinct functions, summarized in the pie chart in Figure 3. The fold change in selected genes based on microarray that we hypothesized were important for mineralization are shown in Table II.

Several of these genes hypothesized to be important for the mineralization process based on level of expression were selected for confirmation by qRT-PCR. The genes that were selected as of interest for the mineralization process were extracellular matrix proteins, the enzymes that modify them, and a sodium phosphate co-transporter.

Bone morphogenic protein 1 (BMP1), a metalloproteinase that activates collagen, several growth factors, and the extracellular matrix protein dentin matrix protein 1 (DMP1) [Reynolds et al., 2000; Ge and Greenspan, 2006] showed increasing gene expression (Fig. 4A) relative to non-mineralizing cultures throughout the culture period, although the increase only reached statistical significance at day 21. DMP-1, a Small Integrin-Binding Ligand N-Glycosylated (SIBLING) protein associated with initiation of mineralization, showed significantly increased expression on days 9–12, just as mineralization commenced (Fig. 4B). The metalloproteinase MMP-13, important for cartilage modification, and believed to be important for preparing the growth plate for vascular invasion [Malemud, 2006] was highest in expression prior to the commencement of mineralization, and then gradually approached equivalence with values in the non-mineralizing cultures (Fig. 4C). Alkaline phosphatase gene expression was significantly elevated relative to the non-mineralizing controls at days 7 and 9, and then gradually decreased so that by day 21 expression was higher in the non-mineralizing controls (Fig. 4D from 3D). The sodium-phosphate co-transporter, NaPi IIb (Fig. 4E) was elevated before mineralization commenced, remained elevated during initial mineralization and was equivalent to the non-mineralizing controls at later time points (days 16 and 21).

Type X collagen, a marker of chondrocyte hypertrophy [Nurminskaya and Linsenmayer, 1996] was elevated relative to non-mineralizing cultures at days 9, 11, and 14 (Fig. 5A), returning to equivalent levels as in the controls by day 16 (Fig. 5A). Types II (Fig. 5B) and I (Fig. 5C) collagen gene showed few significant changes associated with mineralization although there were temporal changes, and type II collagen gene expression was significantly elevated in the mineralizing cultures at day 12. Similarly, bone sialoprotein (BSP) expression was also elevated at day 16 (Fig. 5D) as was osteocalcin (OCN) expression (Fig. 5E). In contrast, osteopontin (Fig. 5F) showed a marked elevation just prior to the time of detectable mineral deposition, and remained elevated through day 16.

DISCUSSION

New information on biomimetic mineralization of the avian growth plate was provided by these analyses, specifically the lack of major variation in collagen I and II gene expression when

mineralizing and non-mineralizing cultures were compared at the same time points, the early upregulation of the osteopontin gene, and the upregulation of the sodium-phosphate co-transporter, the protease BMP1, and one of its substrates, DMP1. Confirmatory information relative to earlier results in the chick system was given for Type X collagen [Boskey et al., 1992b], and, in other growth plate systems by the up-regulation of the matrix metalloproteinases [Wang et al., 2004b]. Similar to other reports [James et al., 2010] we found correspondence between our microarray and qRT-PCR data. In this light it is interesting to note that our microarray data, similar to other studies describing gene regulation in the hypertrophic zone of the growth plate (not mineralization), also reported up-regulation of type I collagen, lysyl oxidase, ADAMTS1, pro col V, and I, BMP2 and 6, ADAM17, calmodulin I, FGFRI, and Annexin I compared to the proliferating zone [Wang et al., 2004b]. Novel among our findings was the upregulation of the Na Pi II b co-transporter during the mineralization process.

The Na Pi IIb cotransporter, upregulated during both the start of mineral deposition and during mineral proliferation, was first isolated from kidney membranes [Debiec et al., 1992], and is the most abundant transporter in murine kidneys [Miyamoto et al., 1997]. Type II co-transporters, in general, play a major role in the regulation of renal Pi reabsorption by dietary Pi and parathyroid hormone, which in turn regulate the endocytosis/exocytosis of the transporters [Takeda et al., 1999]. In chondrocytes, Mansfield et al. [2001] using terminally differentiated hypertrophic chick chondrocytes and a competitive inhibitor of the transporter, phosphonoformic acid, blocked anion-induced cell death and prevented an increase in the cell Pi content. They suggested that at the mineralization front, cell death was linked directly to the elevation in environmental anion concentration and the concomitant rise in intracellular Pi levels. In a murine mesenchymal cell line Denison et al. [2009] showed this NaPi co-transporter was involved in chondrocyte differentiation. The NaPi IIb co-transporter is also important in other mineralized tissue forming cells. Both rat odontoblasts [Lundquist et al., 2002] and osteoblast-like cell lines [Lundquist et al., 2007] were shown to be dependent for mineralization on the activity of a NaPi-II cotransporter. In the rat, NaPi II transporter expression is both regulated by Pi in a manner consistent with their playing a role in transcellular Pi flux during mineralization and appears to be involved in the mineralization process. The cellular Pi is presumably used for ATP synthesis, which in turn may promote apoptosis of growth plate chondrocytes through activation of the caspase signal pathways [Roy et al., 2010]. Extracellular matrix vesicles also use phosphate transport to accelerate mineralization, demonstrated by inhibiting phosphate transport in chick growth plate cultures [Wu et al., 2002].

It is also important to comment on the up-regulation of the BMP1 gene with increasing mineral proliferation. BMP1 (bone morphogenetic protein 1 or tolloid-like metalloproteinase) [Ge and Greenspan, 2006] is the enzyme responsible for the cleavage of dentin matrix protein-1, DMP-1, among other non-collagenous and collagenous matrix proteins into their component fragments [Steiglitz et al., 2004; Von Marschall and Fisher, 2008]. Although relative upregulation of DMP-1 expression preceded BMP1 elevation in the mineralizing vs. non-mineralizing cultures, BMP-1 is likely to be upregulated in both culture systems, as it is needed for processing collagen and lysyl-oxidase, as well as proteins more associated with mineralization. Animals with a mutated BMP1 cleavage site have decreased mineralization (Qin, personal communication), as do hypophosphatemic DMP-1 knockout mice [Liu et al., 2008]. This suggests that BMP1 is needed to prepare the newly formed matrix for mineralization.

The expression of osteocalcin genes were also significantly enhanced after mineralization had commenced, during the stage when mineral crystals are proliferating most actively. Both of these genes have been associated with the presence of mineral in other systems

[Owen et al., 1991]; their expressions are parallel in other systems [Wang et al., 2004a], however, their precise roles in cartilage calcification are not well defined.

In conclusion, these studies enhanced our understanding of the mechanism that allows the chick limb-bud mesenchymal cell cultures to form physiologic hydroxyapatite deposits, and suggest how the process can be further examined (by knockdown or overexpression experiments) to gain additional insights into mineralization mechanisms, pointing out the potential roles of specific enzymes and phosphate transport systems in the mineralization process. Limitations of the study include the fact that avian and mammalian growth plate mineralization may not be identical and the lack of information on many of the up- and down-regulated genes on the chick microarray. None the less this study indicated a list of factors and extracellular matrix components that can be further studied to define their roles in the mineralization process.

Acknowledgments

Grant sponsor: NIH; Grant number: AR037661.

References

- Ahrens PB, Solursh M, Reiter RS. Stage-related capacity for limb chondrogenesis in cell culture. *Dev Biol.* 1977; 60:69–82. [PubMed: 198274]
- Belluccio D, Wilson R, Thornton DJ, Wallis TP, Gorman JJ, Bateman JF. Proteomic analysis of mouse growth plate cartilage. *Proteomics.* 2006; 6:6549–6553. [PubMed: 17163436]
- Binderman I, Greene RM, Pennypacker JP. Calcification of differentiating skeletal mesenchyme in vitro. *Science.* 1979; 206:222–225. [PubMed: 482937]
- Boskey AL, Stiner D, Doty SB, Binderman I. Requirement of vitamin C for cartilage calcification in a differentiating chick limb-bud mesenchymal cell culture. *Bone.* 1991; 12:277–282. [PubMed: 1793679]
- Boskey AL, Camacho NP, Mendelsohn R, Doty SB, Binderman I. FT-IR microscopic mappings of early mineralization in chick limb bud mesenchymal cell cultures. *Calcif Tissue Int.* 1992a; 51:443–448. [PubMed: 1451012]
- Boskey AL, Stiner D, Doty SB, Binderman I, Leboy P. Studies of mineralization in tissue culture: optimal conditions for cartilage calcification. *Bone Miner.* 1992b; 16:11–36. [PubMed: 1371424]
- Boskey AL, Doty SB, Stiner D, Binderman I. Viable cells are a requirement for in vitro cartilage calcification. *Calcif Tissue Int.* 1996; 58:177–185. [PubMed: 8852573]
- Boskey AL, Stiner D, Binderman I, Doty SB. Effects of proteoglycan modification on mineral formation in a differentiating chick limb-bud mesenchymal cell culture system. *J Cell Biochem.* 1997; 64:632–643. [PubMed: 9093912]
- Boskey AL, Stiner D, Binderman I, Doty SB. Type I collagen influences cartilage calcification: an immunoblocking study in differentiating chick limb-bud mesenchymal cell cultures. *J Cell Biochem.* 2000; 79:89–102. [PubMed: 10906758]
- Boskey AL, Doty SB, Kudryashov V, Mayer-Kuckuk P, Roy R, Binderman I. Modulation of extracellular matrix protein phosphorylation alters mineralization in differentiating chick limb-bud mesenchymal cell micro-mass cultures. *Bone.* 2008; 42:1061–1071. [PubMed: 18396125]
- Cameron TL, Belluccio D, Farlier PG, Brachvogel B, Bateman JF. Global comparative transcriptome analysis of cartilage formation in vivo. *BioMed Central Developmental Biology.* 2009; 9:20. [PubMed: 19272164]
- Dean RB, Dixon WJ. Simplified Statistics for Small Numbers of Observations. *Anal Chem.* 1951; 23:636–6638.
- Debiec H, Lorenc R, Ronco PM. Reconstitution and characterization of a Na⁺/Pi co-transporter protein from rabbit kidney brush-border membranes. *Biochem J.* 1992; 286:97–102. [PubMed: 1520289]

- Denison TA, Koch CF, Shapiro IM, Schwartz Z, Boyan BD. Inorganic phosphate modulates responsiveness to 24,25(OH)2D3 in chondrogenic ATDC5 cells. *J Cell Biochem*. 2009; 107:155–162. [PubMed: 19288498]
- Ge G, Greenspan DS. Developmental roles of the BMP1/TLD metalloproteinases. *Birth Defects Res C Embryo Today*. 2006; 78:47–68. [PubMed: 16622848]
- Hamburger V, Hamilton HL. A series of normal stages in the development of the chick embryo. *J Morphol*. 1951; 88:49–92.
- James CG, Stanton L-A, Agoston H, Ulici V, Underhill TM, Beier F. Genome-wide analysis of gene expression during mouse endochondral ossification. *PLOS One*. 2010; 5:e8693. [PubMed: 20084171]
- Khan SN, Solaris J, Ramsey KE, Yang X, Bostrom MP, Stephan D, Daluiski A. Identification of novel gene expression in healing fracture callus tissue by DNA microarray. *HSS J*. 2008; 4:149–160. [PubMed: 18752025]
- Kuroda S, Virdi AS, Dai Y, Shott S, Sumner DR. Patterns and localization of gene expression during intramembranous bone regeneration in the rat femoral marrow ablation model. *Calcif Tissue Int*. 2005; 77:212–225. [PubMed: 16193234]
- Liu S, Zhou J, Tang W, Menard R, Feng JQ, Quarles LD. Pathogenic role of Fgf23 in Dmp1-null mice. *Am J Physiol Endocrinol Metab*. 2008; 295:E254–E261. [PubMed: 18559986]
- Lundquist P, Ritchie HH, Moore K, Lundgren T, Linde A. Phosphate and calcium uptake by rat odontoblast-like MRPC-1 cells concomitant with mineralization. *J Bone Miner Res*. 2002; 17:1801–1813. [PubMed: 12369784]
- Lundquist P, Murer H, Biber J. Type II Na+-Pi cotransporters in osteoblast mineral formation: regulation by inorganic phosphate. *Cell Physiol Biochem*. 2007; 19:43–56. [PubMed: 17310099]
- Malemud CJ. Matrix metalloproteinases: role in skeletal development and growth plate disorders. *Front Biosci*. 2006; 11:1702–1715. [PubMed: 16368549]
- Mansfield K, Teixeira CC, Adams CS, Shapiro IM. Phosphate ions mediate chondrocyte apoptosis through a plasma membrane transporter mechanism. *Bone*. 2001; 28:1–8. [PubMed: 11165936]
- Miyamoto K, Segawa H, Morita K, Nii T, Tatsumi S, Taketani Y, Takeda E. Relative contributions of Na+-dependent phosphate co-transporters to phosphate transport in mouse kidney: RNase H-mediated hybrid depletion analysis. *Biochem J*. 1997; 327:735–739. [PubMed: 9581550]
- Nurminskaya M, Linsenmayer TF. Identification and characterization of up-regulated genes during chondrocyte hypertrophy. *Dev Dyn*. 1996; 206:260–271. [PubMed: 8896982]
- Owen TA, Aronow MS, Barone LM, Bettencourt B, Stein GS, Lian JB. Pleiotropic effects of vitamin D on osteoblast gene expression are related to the proliferative and differentiated state of the bone cell phenotype: dependency upon basal levels of gene expression, duration of exposure, and bone matrix competency in normal rat osteoblast cultures. *Endocrinology*. 1991; 128:1496–1504. [PubMed: 1999168]
- Reynolds SD, Zhang D, Puzas JE, O'Keefe RJ, Rosier RN, Reynolds PR. Cloning of the chick BMP1/Tolloid cDNA and expression in skeletal tissues. *Gene*. 2000; 248:233–243. [PubMed: 10806368]
- Roy R, Kudryashov V, Binderman I, Boskey AL. The role of apoptosis in mineralizing murine vs. avian micromass culture systems. *J Cell Biochem*. 2010; 111:653–658. [PubMed: 20589756]
- Rundle CH, Wang H, Yu H, Chadwick RB, Davis EI, Wergedal JE, Lau KH, Mohan S, Ryaby JT, Baylink DJ. Microarray analysis of gene expression during the inflammation and endochondral bone formation stages of rat femur fracture repair. *Bone*. 2006; 38:521–5529. [PubMed: 16321582]
- Steiglitz BM, Ayala M, Narayanan K, George A, Greenspan DS. Bone morphogenetic protein-1/Tolloid-like proteinases process dentin matrix protein-1. *J Biol Chem*. 2004; 279:980–986. [PubMed: 14578349]
- Takeda E, Taketani Y, Morita K, Miyamoto K. Sodium-dependent phosphate co-transporters. *Int J Biochem Cell Biol*. 1999; 31:377–381. [PubMed: 10224663]
- Ulici V, James CG, Hoenselaar KD, Beier F. Regulation of gene expression by PI3K in mouse growth plate chondrocytes. *PLOS One*. 2010; 5(3):8866.
- Von Marschall Z, Fisher LW. Dentin matrix protein-1 isoforms promote differential cell attachment and migration. *J Biol Chem*. 2008; 283:32730–32740. [PubMed: 18819913]

- Wang C, Duan Y, Markovic B, Barbara J, Howlett CR, Zhang X, Zreiqat H. Phenotypic expression of bone-related genes in osteoblasts grown on calcium phosphate ceramics with different phase compositions. *Biomaterials*. 2004a; 25:2507–2514. [PubMed: 14751735]
- Wang Y, Middleton F, Horton JA, Reichel L, Farnum CE, Damron TA. Microarray analysis of proliferative and hypertrophic growth plate zones identifies differentiation markers and signal pathways. *Bone*. 2004b; 35:1273–1293. [PubMed: 15589209]
- Wang K, Vishwanath P, Eichler GS, Al-Sebaei MO, Edgar CM, Einhorn TA, Smith TF, Gerstenfeld LC. Analysis of fracture healing by large-scale transcriptional profile identified temporal relationships between metalloproteinase and ADAMTS mRNA expression. *Matrix Biol*. 2006; 25:271–281. [PubMed: 16584876]
- Wu LN, Guo Y, Genge BR, Ishikawa Y, Wuthier RE. Transport of inorganic phosphate in primary cultures of chondrocytes isolated from the tibial growth plate of normal adolescent chickens. *J Cell Biochem*. 2002; 6:475–489. [PubMed: 12210754]
- Xiao Z, Camalier CE, Nagashima K, Chan KC, Lucas DA, de la Cruz MJ, Gignac M, Lockett S, Issaq HJ, Veenstra TD, Conrads TP, Beck GR Jr. Analysis of the extracellular matrix vesicle proteome in mineralizing osteoblasts. *J Cell Physiol*. 2007; 210:325–335. [PubMed: 17096383]
- Zhang M, Pritchard MR, Middleton FA, Horton JA, Damron TA. Microarray analysis of perichondral and reserve growth plate zones indentifies differential gene expressions and signal pathways. *Bone*. 2008; 43:511–520. [PubMed: 18579462]





Fig. 1.
Schematic showing stages of matrix and mineral development in the chick limb-bud micromass culture system with typical ^{45}Ca uptake data for mineralizing (dark bars) and non-mineralizing (light bars) as % of day 21 values in mineralizing cultures. The titles in boxes define the stages; the dark arrows indicate the times sampled for micro-array analysis, and the small arrows indicate the times mineralized and non-mineralized cultures were analyzed by qRT-PCR.

**Fig. 2.**

K cluster analyses indicating variation in gene expression with mineralization and time. The genes listed under each figure are those which have been implicated in other studies to the progression of endochondral ossification and/or growth plate mineralization.






Fig. 3.

Distribution of up- and down- regulated genes at day 11 by category.

**Fig. 4.**

Relative expression of A) BMP1, B) DMP-1, C) MMP-13, D) alkaline phosphatase, and E) NaPi IIb. Values are fold change relative to non-mineralizing controls, mean \pm SD, n = 3 or 4. * $P<0.05$ relative to equivalence (constant = 1).

**Fig. 5.**

Relative expression of collagens A) type X, b) type II, c) type I, and extracellular matrix proteins D) bone sialoprotein, E) osteocalcin and F) osteopontin. Values are fold change relative to non-mineralizing controls, mean \pm SD, n = 3 or 4. *P<0.05 relative to equivalence (constant =1).

TABLE I

Primers Used in this Study

Gene	Forward primer	Reverse primer
<i>BMP1</i>	QIAGEN CAT#: QT01491812	
<i>NaPi IIb</i>	CTGGATGCACTCCCTAGAGC	TTATCTTGGCACCCCTCTG
<i>MMP13</i>	CAACCCAAAACATCCAAAAC	CCATTCATAGCCCCAACCTTC
<i>DMP1</i>	QIAGEN CAT#: QT01138354	
<i>Col X</i>	AGTGCTGTATTGATCTCATGGA	TCAGAGGAATAGAGACCATTGGATT
<i>Col II</i>	GACCTCGTGGTGACAAAGGT	CATGCCGTTAGAGCCATCTT
<i>Col I</i>	GCCGTCACCTCAGACTTAGC	TTTGTCCTTGGGTTCTTG
<i>OPN</i>	CCAGCTCTGAAGAAAAATAC	CTAGGAATGTCAGGAAAGTC
<i>OCN</i>	TCGGCGCGCTGCTCACATTCA	TGGCGGTGGGAGATGAAGGCTTA

TABLE II

Microarray Results

Gene	Day 11	Day 17
Extracellular matrix molecules		
DMP1	6.3	2.5
Ovocleidin 116 (MEPE)	5.3	2.4
SPP24 (SPP2)	4.5	0.24
Type IIA collagen precursor		5.1
Type XIV collagen		2.5
Type XII collagen		0.48
Osteoglycin		24
Phosphate Regulation		
Na Pi II Co-transporter	3.1	2.6
Phosphatases (5 distinct genes)	4–12x	2–4
Spingosine phosphatase (also called Phospho1)		8.2
Enzymes (Proteinases & Aggrecanases)		
BMP1	2.6	3.1
MMP15	3	0.5
Stromelysin 3		2.1
MT3-MMP		0.36
ADAMTS-2 (procollagen III N-proteinase)		0.2